Dw	
\boldsymbol{n}	Patient Name

П	_	+	^	
ப	а	L	C	

Prescription for Pain Self-Management

Acetaminophen (i.e., Tylenol®). Take two 325 mg tablets by mouth every six hours as needed for pain	Pain relieving cream or rub (i.e., IcyHot®, Arnicare Gel®): use as directed on the package		
Acetaminophen (i.e., Tylenol®). Take two 500 mg tablets by mouth every six hours as needed for pain (not to exceed 4000mg)	Apply ice or cold pack for 15-20 minutes, 3 times a day		
Ibuprofen (i.e., Advil® or Motrin®). Take two 200 mg tablets by mouth every six hours as needed for pain	Apply heat pack for 15-20 minutes, 3 times a day Other:		
Combination therapy. Take one 500 mg tablet of extra strength acetaminophen AND one 200 mg tablet of ibuprofen by mouth every six hours as needed for pain			
Alternating therapy. Take two 325 mg tablets of acetaminophen by mouth NOW, THEN take two 200 mg tablets of ibuprofen four hours later	If not improved in days or if new symptoms occur, call or return to the office for a recheck.		
Example: acetaminophen at 8 am, ibuprofen at 12 pm, acetaminophen at 4 pm, ibuprofen at 8 pm, continue as needed	* Note : Tylenol® and store brands of acetaminophen are the same and work equally well. Advil®, Motrin® and store brands of ibuprofen are the same and work equally well. Maximum dose : NSAIDS 3200mg and Acetaminophen		
You have NOT been prescribed a narcotic (opioid) medication because the risks outweigh the benefits for your condition.	4000mg in a 24-hour period.		
* Opioids have a risk for addiction/misuse	Prescriber:		



